

OPTIMAL CONTROL OF A LARGE DAM, TAKING INTO ACCOUNT THE WATER COSTS

VYACHESLAV M. ABRAMOV

ABSTRACT. Consider a dam model, L^{upper} and L^{lower} are upper and, respectively, lower levels, $L = L^{upper} - L^{lower}$ is large and if the level of water is between these bounds, then the dam is said to be in a normal state. Passage across lower or upper levels leads to damage. Let $J_1 = j_1 L$ and $J_2 = j_2 L$ denote the damage costs per time unit of crossing the lower and, correspondingly, upper level where j_1 and j_2 are given real constants. It is assumed that input stream of water is described by a Poisson process, while the output stream is state dependent. Let L_t denote the level of water in time t , and c_{L_t} denote the water cost at level L_t ($L^{lower} < L_t \leq L^{upper}$). Assuming that $p_1 = \lim_{t \rightarrow \infty} \mathbf{P}\{L_t = L^{lower}\}$, $p_2 = \lim_{t \rightarrow \infty} \mathbf{P}\{L_t > L^{upper}\}$ and $q_i = \lim_{t \rightarrow \infty} \mathbf{P}\{L_t = i\}$ ($L^{lower} < i \leq L^{upper}$) exist, the aim of the paper is to choose the parameters of an output stream (specifically defined in the paper) minimizing the long-run expenses

$$J = p_1 J_1 + p_2 J_2 + \sum_{i=L^{lower}+1}^{L^{upper}} q_i c_i.$$

1. INTRODUCTION

A large dam is defined by the parameters L^{lower} and L^{upper} , which are, respectively, the lower and upper levels of the dam. If the current level is between these bounds, the dam is assumed to be in a normal state. The difference $L = L^{upper} - L^{lower}$ is large, and this is the reason for calling the dam *large*. This property enables us to use asymptotic analysis as $L \rightarrow \infty$ and solve easier different problems of optimal control than we would were than the dam is not large.

Let L_t denote the water level in time t . If $L^{lower} < L_t \leq L^{upper}$, then the state of the dam is called *normal*. Passage across lower or upper level leads to damage. The costs per time unit of this damage is $J_1 = j_1 L$ and $J_2 = j_2 L$ for lower and upper levels correspondingly, where j_1 and j_2 are given real constants. The water inflow is described by the Poisson process with rate λ . In practice, this means that the arrival of water units is registered by counter at random instants $t_1, t_2, \dots$, and the times between consecutive instants are mutually independent and exponentially distributed with parameter λ .

The outflow of water is state-dependent as follows. If the level of water is between L^{lower} and L^{upper} , then an interval between departures of units of water (inverse output flow) has the probability distribution function $B_1(x)$. If level of water exceeds L^{upper} , then an inverse output flow has the probability distribution function $B_2(x)$. The probability distribution function $B_2(x)$ is assumed to obey the condition

1991 *Mathematics Subject Classification.* 60K30, 40E05, 90B05, 60K25.

Key words and phrases. Dam, State-dependent queue, Asymptotic analysis, Control problem.

$\int_0^\infty x dB_2(x) < \frac{1}{\lambda}$. If the level of water is L^{lower} exactly, then output of water is frozen and it resumes again as soon as the level of water exceeds the level L^{lower} .

Let c_{L_t} denote the cost of water at level L_t . The sequence c_i is assumed to be positive and non-increasing. The problem of the present paper is to choose the parameter $\int_0^\infty x dB_1(x)$ of the dam in the normal state minimizing the objective function

$$(1.1) \quad J = p_1 J_1 + p_2 J_2 + \sum_{i=L^{lower}+1}^{L^{upper}} c_i q_i,$$

where

$$(1.2) \quad p_1 = \lim_{t \rightarrow \infty} \mathbf{P}\{L_t = L^{lower}\},$$

$$(1.3) \quad p_2 = \lim_{t \rightarrow \infty} \mathbf{P}\{L_t > L^{upper}\},$$

$$(1.4) \quad q_i = \lim_{t \rightarrow \infty} \mathbf{P}\{L_t = L^{lower} + i\}, \quad i = 1, 2, \dots, L.$$

In the queueing formulation, the level L^{lower} is identified with an empty queue, and the dam model is the following queueing system with service depending on queue-length. If immediately before a service begins the queue-length exceeds the level L , then the customer is served by the probability distribution $B_2(x)$. Otherwise, the service time distribution is $B_1(x)$. The value p_1 is the stationary probability of empty system, the value p_2 is the stationary probability that a customer is served by probability distribution $B_2(x)$, and q_i , $i = 1, 2, \dots, L$, are the stationary probabilities of the queue-length process, so $p_1 + p_2 + \sum_{i=1}^L q_i = 1$. (For the described queueing system, the right-hand side limits in relations (1.2)-(1.4) do exist.)

In our study, the parameter L increases indefinitely, and we deal with the series of queueing systems. The parameters above, such as p_1 , p_2 , J_1 , J_2 as well as other parameters are functions of L . The argument L will be often omitted in these functions.

Similarly to [5], it is assumed that the input parameter λ and probability distribution function $B_2(x)$ are given, while the appropriate probability function $B_1(x)$ should be chosen from the specified parametric family of functions $B_1(x, C)$. (Actually, we deal with the family of probability distributions $B_1(x)$ depending on two parameters δ and L in series. Then the parametric family of distributions $B_1(x, C)$ is described by the family of possible limits of δL as $\delta \rightarrow 0$ and $L \rightarrow \infty$.)

The outflow rate associated with the probability distribution function $B_1(x)$ should be chosen such that the minimum of the objective function of (1.1) is associated with the choice of the parameter C resulting in the choice of the corresponding probability distribution function $B_1(x, C)$.

The more particular problem, where the objective function has the form $J = p_1 J_1 + p_2 J_2$ (i.e. the water costs are not taken into account), has been studied in Abramov [5]. In this case the solution to the control problem is unique and has one of the following three forms. Denote $\rho_2 = \lambda \int_0^\infty x dB_2(x)$ and $\rho_1 = \rho_1(C) = \lambda \int_0^\infty x dB_1(x, C)$. Then, in the case $j_1 = j_2 \frac{\rho_2}{1-\rho_2}$, the optimal solution is $\rho_1 = 1$. In the case $j_1 > j_2 \frac{\rho_2}{1-\rho_2}$, the optimal solution has the form $\rho_1 = 1 + \delta$, where $\delta(L)$ is a small positive parameter, and $\delta(L)L \rightarrow C$ as $L \rightarrow \infty$. In the case $j_1 < j_2 \frac{\rho_2}{1-\rho_2}$, the optimal strategy has the form $\rho_1 = 1 - \delta$, and $\delta(L)L \rightarrow C$ as $L \rightarrow \infty$. The parameter C is a unique solution of a specific minimization problem precisely formulated in [5].

It has been also shown in [5] that the solution to the control problem is insensitive to the type of probability distributions $B_1(x)$ and $B_2(x)$. Specifically, it is expressed via the first moment of $B_2(x)$ and the first two moments of $B_1(x)$.

Following [5], we use the notation $\rho_{1,l} = \lambda^l \int_0^\infty x^l dB_1(x)$, $l = 2, 3$. The existence of a moment of the order corresponding to $\rho_{1,l}$ will be specially assumed in formulations of statements corresponding to case studies.

The problem studied in the present paper substantially distinguishes from that studied in [5]. Although the both control problem of the present paper and [5] are closely related, the new components of the present problem change the problem substantially. The problem of the present paper requires a much deepen and delicate analysis (for example, it is demonstrated in the next section that the asymptotic methods of the earlier paper [5] do not longer work here, and one should use more delicate techniques instead), and the main results on optimal control policies are deepen as well.

Essential difficulty of the control problem in the present formulation is to prove a uniqueness of the optimal solution, while in the case of the particular problem of [5], the uniqueness of the solution follows automatically from the explicit representations of the functionals obtained there.

It is assumed in the present paper that c_i is a non-increasing sequence. If the cost sequence c_i were an arbitrary bounded sequence, then a richer class of possible cases could be studied. However, in the case of arbitrary cost sequence, the solution need not be unique, and arbitrary costs c_i , say increasing in i , seem not to be useful and, therefore, are not considered here. The practical applications of the results obtained in the paper are not restricted by the area of water research. The results of the present paper are meaningful for many specific inventory and storage problems as well. Moreover, the model considered in this paper describes more realistically a production/realization process of large warehouses rather than an inflow/outflow process of large dams, because in many cases of a water inflow the seasonality is important. Nevertheless, similarly to [5], the problem formulation in this paper is given in terms of water research.

More realistic models arising in practice assume that the probability distribution function $B_1(x)$ should also depend on i , i.e have representation $B_{1,i}(x)$. The model of the present paper, where $B_1(x)$ is the same for all i , under appropriate additional information can approximate those more general models. Namely, one can suppose the stationary service time distribution $B_1(x)$ has the representation $B_1(x) = \sum_{i=1}^L q_i B_{1,i}(x)$ (q_i , $i = 1, 2, \dots, L$ are the state probabilities), and the solution to the control problem for $B_1(x)$ enables us to find then the approximate solutions to the control problem for $B_{1,i}(x)$, $i = 1, 2, \dots, L$ by using the Bayes rule.

Similarly to the solution of the control problem of [5], the solution of the present problem with extended criteria (1.1) is related to the same class of solutions as in [5]. That is, it must be either $\rho_1 = 1$ or one of two limits of $\rho_1 = 1 + \delta$, $\rho_1 = 1 - \delta$ for positive small vanishing δ as L increases indefinitely, and $L\delta \rightarrow C$. The reason for this is that the penalties upon reaching upper or lower level are of order $O(L)$ (i.e. increase to infinity as $L \rightarrow \infty$), while the water costs are assumed to be bounded as L tends to infinity, and although the water costs affect the solution of the control problem, this influence remains in the framework of the same class of solutions mentioned above.

The following new questions are of special interest here.

1. What is the structure of an optimal solution? Is an optimal solution unique?

The answer to these questions is the main result of the paper. The questions are answered by Theorem 3.3. We prove that a solution to the control problem does exist and unique, however there are some additional mild assumptions related to the class of probability distributions $\{B_1(x)\}$. The proof of the existence and uniqueness of a solution is based on special techniques of Mathematical Analysis. Specifically, we use the known techniques of majorization inequalities [7], [8] in order to prove the monotonicity of specified functions. This property of monotonicity is then used to prove a uniqueness of a solution.

2. Under what relation between j_1 , j_2 , ρ_2 , c_i (and maybe other parameters of the model) the optimal strategy is $\rho_1 = 1$?

If the water costs are not taken into account, then the condition for $\rho_1 = 1$ is $j_1 = j_2 \frac{\rho_2}{1-\rho_2}$. This result has been proved in [5]. It has a simple intuitive explanation and is a consequence of the well-known property of the stream of lost calls during a busy period of M/GI/1/n queues, under the assumption that the expected interarrival and service times are equal (see Abramov [1] as well as Righter [10] or Wolff [16]). Taking into account the structure of water costs generally changes this condition for the aforementioned optimal solution $\rho_1 = 1$. We prove that the optimal solution $\rho_1 = 1$ is achieved under the condition $j_1 \leq j_2 \frac{\rho_2}{1-\rho_2}$, and the equality in this relation holds if and only if the water costs are the same at all levels of water. This result is only the partial answer to the question. More exact answers can be obtained in particular cases, and one of them is the case of linearly decreasing costs as the level of water increases (for brevity, this case is called *linear costs*). In the case of linear costs we derive more exact and useful representations, which enable us to calculate numerically the relation between j_1 and j_2 to have finally the optimal solution $\rho_1 = 1$. The relevant numerical results are provided for special values of the parameters of the model.

The rest of the paper is organized as follows. In Section 2, the asymptotic behavior of the stationary probabilities is studied. This section is structured into four subsections. In Section 2.1, some necessary results related to asymptotic properties of characteristic of the state-dependent queueing system are recalled for modelling the behavior of a large dam. In Section 2.2, elementary asymptotic properties of the state probabilities in the case $\rho_1 = 1$ are established. The analysis of this case is based on application of a Tauberian theorem of Postnikov [9] for the convolution type recurrence relation. In Section 2.3 the behavior of the state probabilities is studied for the case $\rho_1 = 1 + \delta$, where positive parameter δ vanishes as L tends to infinity. The analysis of this case uses known results on asymptotic behavior of characteristics of the state-dependent queueing system obtained in Section 2.1. Delicate asymptotic analysis is provided in Section 2.4 for the case $\rho_1 = 1 - \delta$, where positive parameter δ vanishes as L tends to infinity. For the analysis of this case we involve the asymptotic result of Willmot [15], which is also used in the present paper. All of the aforementioned asymptotic results of Section 2 are used to establish asymptotic properties of special functionals. Then we solve the problem of minimization of these functionals. In Section 3, we solve the control problem. We prove existence and uniqueness of the solution under mild assumptions on the class of probability distributions $\{B_1(x)\}$. We establish a structure of the solution and completely answer the question 1 posed in the introduction. In section 4, the case of linear costs is studied. We establish explicit representations for functionals of

the control problem and provide numerical answer to the question 2 posed in the introduction.

2. STATIONARY PROBABILITIES OF THE STATE-DEPENDENT QUEUEING SYSTEM AND THEIR ASYMPTOTIC BEHAVIOR

In this section, the explicit expressions are derived for the stationary probabilities, and their asymptotic behavior is studied. These results will be used in our further findings of the optimal solution.

2.1. Preliminaries. We first recall some results of [5] and then develop them in order to obtain the explicit representations for the stationary probabilities that will be required in our further analysis. Let T_L , ν_L , I_L denote correspondingly a busy period, the number of served customers during a busy period, and an idle period. Let $\nu_L^{(1)}$ and $\nu_L^{(2)}$ denote the number of customers served during a busy period by probability distribution functions $B_1(x)$ and $B_2(x)$ correspondingly, and let $T_L^{(1)}$ and $T_L^{(2)}$ denote the time spent for the service of customers during a busy period by the probability distribution functions $B_1(x)$ and $B_2(x)$ correspondingly.

It was shown in [5] that the probabilities p_1 and p_2 can be expressed explicitly via $\mathbf{E}\nu_L^{(1)}$, ρ_1 and ρ_2 only. Namely,

$$(2.1) \quad p_1 = \frac{1 - \rho_2}{1 + (\rho_1 - \rho_2)\mathbf{E}\nu_L^{(1)}},$$

and

$$(2.2) \quad p_2 = \frac{\rho_2 + \rho_2(\rho_1 - 1)\mathbf{E}\nu_L^{(1)}}{1 + (\rho_1 - \rho_2)\mathbf{E}\nu_L^{(1)}}.$$

Such kind of representations is convenient, because $\mathbf{E}\nu_L^{(1)}$ satisfies the convolution type recurrence relation

$$(2.3) \quad \mathbf{E}\nu_L^{(1)} = \sum_{j=0}^L \mathbf{E}\nu_{L-j+1}^{(1)} \int_0^\infty e^{-\lambda x} \frac{(\lambda x)^j}{j!} dB_1(x), \quad \mathbf{E}\nu_0^{(1)} = 1,$$

where $\mathbf{E}\nu_n^{(1)}$ denotes the expectation of the number of served customers during a busy period of the M/GI/1/n queue ($n=0,1,\dots$). Recurrence relation (2.3) is in turn a special case of the recurrence relations

$$(2.4) \quad Q_n = \sum_{j=0}^n Q_{n-j+1} r_j,$$

with $r_0 > 0$, $r_j \geq 0$ for all $j \geq 1$, and $r_0 + r_1 + \dots = 1$. The detailed theory of these recurrence relations can be found in Takács [14]. Asymptotic behavior of Q_n as $n \rightarrow \infty$ has been studied by Takács [14] and Postnikov [9].

The stationary probabilities q_i can be obtained from the renewal arguments (e.g. Ross [11]). Namely, for $i = 1, 2, \dots, L$ we have

$$(2.5) \quad q_i = \frac{\mathbf{E}T_i^{(1)} - \mathbf{E}T_{i-1}^{(1)}}{\mathbf{E}T_L + \mathbf{E}I_L},$$

where $\mathbf{E}T_i^{(1)}$ denotes the expectation of a busy period of the M/GI/1/ i queue ($i=0,1,\dots$). The probabilities given by (2.5) can be also written

$$(2.6) \quad q_i = \rho_1 \frac{\mathbf{E}\nu_i^{(1)} - \mathbf{E}\nu_{i-1}^{(1)}}{\mathbf{E}\nu_L}.$$

Indeed, from Wald's equation [6], p.384 we have $\mathbf{E}T_i^{(k)} = \frac{\rho_k}{\lambda} \mathbf{E}\nu_i^{(k)}$, $k = 1, 2$, and therefore the numerator of (2.5) is rewritten

$$(2.7) \quad \mathbf{E}T_i^{(1)} - \mathbf{E}T_{i-1}^{(1)} = \frac{\rho_1}{\lambda} \left(\mathbf{E}\nu_i^{(1)} - \mathbf{E}\nu_{i-1}^{(1)} \right).$$

Next, using the fact that the number of arrivals during a busy cycle coincides with the number of served customers during a busy period, from Wald's equation [6], p.384 we obtain

$$(2.8) \quad \lambda \mathbf{E}T_L + \lambda \mathbf{E}I_L = \mathbf{E}\nu_L.$$

Therefore, (2.6) is the consequence of (2.7) and (2.8).

On the other hand, (2.8) can be rewritten

$$(2.9) \quad \begin{aligned} \lambda \mathbf{E}T_L + \lambda \mathbf{E}I_L &= \lambda \mathbf{E}T_L + 1 \\ &= \lambda \left(\mathbf{E}T_L^{(1)} + \mathbf{E}T_L^{(2)} \right) \\ &= \rho_1 \mathbf{E}\nu_L^{(1)} + \rho_2 \mathbf{E}\nu_L^{(2)} + 1. \end{aligned}$$

From (2.8) and (2.9) we have the equation

$$\mathbf{E}\nu_L^{(1)} + \mathbf{E}\nu_L^{(2)} = \rho_1 \mathbf{E}\nu_L^{(1)} + \rho_2 \mathbf{E}\nu_L^{(2)} + 1,$$

resulting in

$$(2.10) \quad \mathbf{E}\nu_L^{(2)} = \frac{1}{1 - \rho_2} - \frac{1 - \rho_1}{1 - \rho_2} \mathbf{E}\nu_L^{(1)}$$

and, consequently by adding to the both sides of (2.10) $\mathbf{E}\nu_L^{(1)}$, we obtain

$$(2.11) \quad \mathbf{E}\nu_L = \frac{1}{1 - \rho_2} + \frac{\rho_1 - \rho_2}{1 - \rho_2} \mathbf{E}\nu_L^{(1)}.$$

Substituting (2.11) for (2.6), we finally obtain

$$(2.12) \quad q_i = \frac{\rho_1(1 - \rho_2)}{1 + (\rho_1 - \rho_2)\mathbf{E}\nu_L^{(1)}} \left(\mathbf{E}\nu_i^{(1)} - \mathbf{E}\nu_{i-1}^{(1)} \right), \quad i = 1, 2, \dots, L.$$

Comparison with (2.1) enables us to rewrite (2.12) in the other form

$$(2.13) \quad q_i = \rho_1 p_1 \left(\mathbf{E}\nu_i^{(1)} - \mathbf{E}\nu_{i-1}^{(1)} \right), \quad i = 1, 2, \dots, L$$

as well.

In order to study the asymptotic behavior of the loss probabilities, let us recall the earlier results on asymptotic behavior of $\mathbf{E}\nu_L^{(1)}$ as L increases indefinitely (e.g. [1], [3], [5]). This asymptotic behavior can be obtained from the results of Takács [14], p.22-23, on asymptotic behavior of recurrence relation (2.4) as $n \rightarrow \infty$. (The formulation of this Takács theorem can be found in [5] as well.)

If $\rho_1 = 1$ and $\rho_{1,2} < \infty$, then

$$(2.14) \quad \lim_{L \rightarrow \infty} \frac{\mathbf{E}\nu_L^{(1)}}{L} = \frac{2}{\rho_{1,2}},$$

and if $\rho_1 > 1$, then

$$(2.15) \quad \lim_{L \rightarrow \infty} \left[\mathbf{E}\nu_L^{(1)} - \frac{1}{\varphi^L(1 + \lambda \widehat{B}'_1(\lambda - \lambda\varphi))} \right] = \frac{1}{1 - \rho_1},$$

where $\widehat{B}_1(s)$ is the Laplace-Stieljes transform of $B_1(x)$, φ is the least in absolute value root of the functional equation $z = \widehat{B}_1(\lambda - \lambda z)$, and $\widehat{B}'_1(s)$ denotes the derivative of $\widehat{B}_1(s)$.

We do not consider the case $\rho_1 < 1$, because it is not meaningful for our analysis. Only relations (2.14) and (2.15) are useful for the asymptotic analysis of the state probabilities, which is provided below.

2.2. The case $\rho_1 = 1$. Asymptotic relation (2.14) is not enough in order to obtain an asymptotic behavior of the state probabilities. As it was mentioned above, this asymptotic relation (2.14) is obtained from the aforementioned Takács theorem [14], which has been also used in [5] for the analysis of the asymptotic behavior of p_1 and p_2 . The asymptotic analysis of the state probabilities is more delicate than that analysis of p_1 and p_2 , and asymptotic relation (2.14) is not enough for this purpose. Therefore, we have to use the more precise asymptotic estimation given by the Tauberian theorem of Postnikov [9] on the asymptotic behavior of recurrence sequence (2.4).

Lemma 2.1. (*Postnikov [9], Sec. 25.*) *Let $\gamma_1 = \sum_{n=1}^{\infty} nr_n = 1$, $\gamma_2 = \sum_{n=2}^{\infty} n(n-1)r_n < \infty$, and $r_0 + r_1 < 1$. Let $Q_0 \neq 0$ be an arbitrarily given initial value of recurrence sequence (2.4). Then as $n \rightarrow \infty$*

$$Q_{n+1} - Q_n = \frac{2Q_0}{\gamma_2} + o(1).$$

From this lemma we obtain the following statement.

Corollary 2.2. *For any $j \geq 0$,*

$$(2.16) \quad \mathbf{E}\nu_{L-j}^{(1)} - \mathbf{E}\nu_{L-j-1}^{(1)} = \frac{2}{\rho_{1,2}} + o(1), \text{ as } L \rightarrow \infty.$$

Proof. In the case where $r_j = \int_0^\infty e^{-\lambda x} \frac{(\lambda x)^j}{j!} dB_1(x)$, $j = 0, 1, \dots$, the Tauberian condition $r_0 + r_1 < 1$ of Lemma 2.1 is always fulfilled, and the result of corollary follows directly from Lemma 2.1. The proof of this fact uses the theory of analytic functions and has been provided in [3] (Theorem 4.6) and [2] (Theorem 3.3). Since this proof is elementary and short it is repeated here again.

We must prove that for *some* $\lambda_0 > 0$ the equality

$$(2.17) \quad \int_0^\infty e^{-\lambda_0 x} (1 + \lambda_0 x) dB_1(x) = 1$$

is not the case. If it is so, then only the inequality

$$\int_0^\infty e^{-\lambda x} (1 + \lambda x) dB_1(x) < 1$$

must hold.

Indeed, $\int_0^\infty e^{-\lambda x} (1 + \lambda x) dB_1(x)$ is an analytic function in λ , and therefore, according to the theorem on the maximum module of an analytic function, equality (2.17) must hold for *all* $\lambda_0 \geq 0$. This means, that (2.17) is valid if and only if

$\int_0^\infty e^{-\lambda_0 x} \frac{(\lambda_0 x)^j}{j!} dB_1(x) = 0$ for all $j \geq 2$ and $\lambda_0 \geq 0$. In this case the Laplace-Stieltjes transform $\widehat{B}_1(\lambda)$ must be a linear function in λ , i.e. $\widehat{B}_1(\lambda) = d_0 + d_1 \lambda$, d_0 and d_1 are some constants. However, since $|\widehat{B}_1(\lambda)| \leq 1$, we have $d_0 = 1$ and $d_1 = 0$. This is a trivial case where $B_1(x)$ is concentrated in point 0, and therefore it is not a probability distribution function having a positive mean. Thus (2.17) is not the case, and the statement of the corollary follows. $\square$

From (2.16) and (2.1), (2.13) we have the following lemma.

Lemma 2.3. *In the case $\rho_1 = 1$ and $\rho_2 < \infty$ for any $j \geq 0$ we have*

$$(2.18) \quad \lim_{L \rightarrow \infty} Lq_{L-j} = 1.$$

Proof. Asymptotic relation (2.18) follows immediately from (2.1), (2.13) and (2.16). $\square$

Combining the result obtained in [5] and Lemma 2.3 we have the asymptotic result for the functional $J = J_1 p_1 + J_2 p_2 + \sum_{i=1}^L q_i c_i$.

Proposition 2.4. *In the case $\rho_1 = 1$ and $\rho_2 < \infty$ we have*

$$(2.19) \quad \lim_{L \rightarrow \infty} J(L) = j_1 \frac{\rho_{1,2}}{2} + j_2 \frac{\rho_2}{1 - \rho_2} \cdot \frac{\rho_{1,2}}{2} + c^*,$$

where

$$c^* = \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{i=1}^L c_i.$$

Proof. The first two term of the right-hand side of (2.19) easily follow from (2.1), (2.2) and asymptotic representation (2.14). The last term of the right-hand side of (2.19) follows from Lemma 2.3, since

$$\lim_{L \rightarrow \infty} \sum_{i=1}^L q_i c_i = \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{i=1}^L c_i = c^*.$$

Notice, that the representation for the first two terms of the right-hand side of (2.19) has a relation to the particular case of the dam model not taking into account the water costs and been obtained in [5]. $\square$

Remark 2.5. The statement of Proposition 2.4 confirms that the class of the possible solutions of the control problem is the same as that for the particular problem studied in [5], since according to Proposition 2.4 the optimal value of the functional J is finite (i.e. does not increase to infinity as L increases indefinitely) and cannot exceed the right-hand side of (2.19).

2.3. The case $\rho_1 = 1 + \delta, \delta > 0$. In the case $\rho_1 = 1 + \delta$ we have the following.

Theorem 2.6. *Assume that $\rho_1 = 1 + \delta, \delta > 0$, and $L\delta \rightarrow C > 0$ as $\delta \rightarrow 0$ and $L \rightarrow \infty$. Assume that $\rho_{1,3}(L)$ is a bounded sequence, and there exists $\tilde{\rho}_{1,2} = \lim_{L \rightarrow \infty} \rho_{1,2}(L)$. Then, for any $j \geq 0$*

$$(2.20) \quad q_{L-j} = \frac{e^{2C/\tilde{\rho}_{1,2}}}{e^{2C/\tilde{\rho}_{1,2}} - 1} \left(1 - \frac{2\delta}{\tilde{\rho}_{1,2}}\right)^j \frac{2\delta}{\tilde{\rho}_{1,2}} + o(\delta).$$

Proof. Expanding first (2.15) for large L , we have

$$(2.21) \quad \mathbf{E}\nu_{L-j}^{(1)} = \frac{\varphi^j}{\varphi^L [1 + \lambda \widehat{B}'_1(\lambda - \lambda\varphi)]} + \frac{1}{1 - \rho_1} + o(1).$$

From (2.21) for large L we have

$$(2.22) \quad \mathbf{E}\nu_{L-j}^{(1)} - \mathbf{E}\nu_{L-j-1}^{(1)} = \frac{(1 - \varphi)\varphi^j}{\varphi^L [1 + \lambda \widehat{B}'_1(\lambda - \lambda\varphi)]} + o(1).$$

Next, under the condition of the theorem we have the following expansion

$$(2.23) \quad \varphi = 1 - \frac{2\delta}{\widetilde{\rho}_{1,2}} + O(\delta^2).$$

(The details of this expansion can be found in [12], p.326 or [4], p.21. See also the proof of a similar fact in Lemma 2.9 below.) Then, taking into account (2.23), we also have

$$(2.24) \quad 1 + \lambda \widehat{B}'_1(\lambda - \lambda\varphi) = \delta + O(\delta^2).$$

Next, taking into account asymptotic expansions (2.23) and (2.24) from (2.21) and (2.22) we have

$$(2.25) \quad \mathbf{E}\nu_L^{(1)} = \frac{e^{2C/\widetilde{\rho}_{1,2}} - 1}{\delta} + O(1),$$

and for any $j = 0, 1, \dots$

$$(2.26) \quad \mathbf{E}\nu_{L-j}^{(1)} - \mathbf{E}\nu_{L-j-1}^{(1)} = e^{2C/\widetilde{\rho}_{1,2}} \left(1 - \frac{2\delta}{\widetilde{\rho}_{1,2}}\right)^j \frac{2}{\widetilde{\rho}_{1,2}} [1 + o(1)].$$

Now, using explicit representation (2.12), for any $j = 0, 1, \dots$ we obtain the desired asymptotic relation

$$q_{L-j} = \frac{e^{2C/\widetilde{\rho}_{1,2}}}{e^{2C/\widetilde{\rho}_{1,2}} - 1} \left(1 - \frac{2\delta}{\widetilde{\rho}_{1,2}}\right)^j \frac{2\delta}{\widetilde{\rho}_{1,2}} [1 + o(1)].$$

The theorem is proved. $\square$

Using the result of [5] we have the following.

Proposition 2.7. *Under the assumptions of Theorem 2.6 denote the objective function J by J^{upper} . We have the following representation*

$$(2.27) \quad J^{upper} = C \left[j_1 \frac{1}{e^{2C/\widetilde{\rho}_{1,2}} - 1} + j_2 \frac{\rho_2 e^{2C/\widetilde{\rho}_{1,2}}}{(1 - \rho_2)(e^{2C/\widetilde{\rho}_{1,2}} - 1)} \right] + c^{upper},$$

where

$$(2.28) \quad c^{upper} = \frac{2C}{\widetilde{\rho}_{1,2}} \cdot \frac{e^{2C/\widetilde{\rho}_{1,2}}}{e^{2C/\widetilde{\rho}_{1,2}} - 1} \lim_{L \rightarrow \infty} \frac{1}{L} \widehat{C}_L \left(1 - \frac{2C}{\widetilde{\rho}_{1,2}L}\right),$$

and $\widehat{C}_L(z) = \sum_{j=0}^{L-1} c_{L-j} z^j$ is a backward generating cost function.

Proof. The representation for the term

$$C \left[j_1 \frac{1}{e^{2C/\widetilde{\rho}_{1,2}} - 1} + j_2 \frac{\rho_2 e^{2C/\widetilde{\rho}_{1,2}}}{(1 - \rho_2)(e^{2C/\widetilde{\rho}_{1,2}} - 1)} \right]$$

is known from [5]. The new term in (2.27), which takes into account the water costs, is c^{upper} . Therefore, keeping in mind representation (2.20), for this term we obtain:

$$c^{upper} = \lim_{L \rightarrow \infty} \sum_{j=0}^{L-1} q_{L-j} c_{L-j} = \lim_{L \rightarrow \infty} \sum_{j=0}^{L-1} c_{L-j} \cdot \frac{e^{2C/\tilde{\rho}_{1,2}}}{e^{2C/\tilde{\rho}_{1,2}} - 1} \left(1 - \frac{2\delta L}{\tilde{\rho}_{1,2}L}\right)^j \frac{2\delta L}{\tilde{\rho}_{1,2}L},$$

and representation (2.28) follows. $\square$

2.4. The case $\rho_1 = 1 - \delta, \delta > 0$. The study of this case is more delicate and based on a special analysis. Our additional assumption here is that the class of probability distribution functions $\{B_1(x)\}$ is given such that there exists a unique root $\tau > 1$ of the equation

$$(2.29) \quad z = \hat{B}(\lambda - \lambda z),$$

and there exists the first derivative $\hat{B}'(\lambda - \lambda\tau)$.

Under the assumption that $\rho_1 < 1$ the root of (2.29) is not necessarily exists. Such type of condition has been considered by Willmot [15] to obtain the asymptotic behavior for high queue-level probabilities in stationary M/GI/1 queues. Denote the stationary probabilities of M/GI/1 queues by $q_i[M/GI/1]$, $i = 0, 1, \dots$. There was shown in [15] that

$$(2.30) \quad q_i[M/GI/1] = \frac{(1 - \rho_1)(1 - \tau)}{\tau^i [1 + \lambda \hat{B}'(\lambda - \lambda\tau)]} [1 + o(1)], \quad \text{as } i \rightarrow \infty.$$

On the other hand, according to the Pollaczek-Khintchine formula (e.g. Takács [13], p. 242) $q_i[M/GI/1]$ can be represented explicitly

$$(2.31) \quad q_i[M/GI/1] = (1 - \rho_1) \left(\mathbf{E}\nu_i^{(1)} - \mathbf{E}\nu_{i-1}^{(1)} \right), \quad i = 1, 2, \dots$$

(Representation (2.31) can be easily checked, since it easily follows from (2.3) that

$$(2.32) \quad \sum_{j=0}^{\infty} \mathbf{E}\nu_j^{(1)} z^j = \frac{\hat{B}_1(\lambda - \lambda z)}{\hat{B}_1(\lambda - \lambda z) - z},$$

and multiplication of the right-hand side of (2.32) by $(1 - \rho_1)(1 - z)$ leads to the well-known Pollaczek-Khintchine formula.)

From (2.30) and (2.31) we also have the following asymptotic proportion. For large L and any $j \geq 0$

$$(2.33) \quad \frac{\mathbf{E}\nu_{L-j}^{(1)} - \mathbf{E}\nu_{L-j-1}^{(1)}}{\mathbf{E}\nu_L^{(1)} - \mathbf{E}\nu_{L-1}^{(1)}} = \tau^j [1 + o(1)].$$

Now we formulate and prove a theorem on asymptotic behavior of stationary probabilities q_i in the case $\rho_1 = 1 - \delta$. In this theorem we assume that the class of probability distributions $\{B_1(x)\}$ is defined according to the above convention. In the case where $\rho_1 = 1 - \delta$, $\delta > 0$ and vanishing δ as L increases indefinitely, this means that there exists $\epsilon_0 > 0$ (small enough) such that for all $0 \leq \epsilon \leq \epsilon_0$, the above family of probability distribution functions $B_{1,\epsilon}(x)$ (depending now on parameter ϵ) satisfies the following properties. Let $\hat{B}_{1,\epsilon}(s)$ denote the Laplace-Stieltjes transform of $B_{1,\epsilon}(x)$. We assume that any $\hat{B}_{1,\epsilon}(s)$ is an analytic function in a small neighborhood of zero, and

$$(2.34) \quad \hat{B}_{1,\epsilon}'(-\lambda\epsilon) < \infty.$$

Property (2.34) is required for the existence of the probabilities q_i . Relation (2.30) contains the term $\widehat{B}'(\lambda - \lambda\tau)$, and for $\tau = 1 + \epsilon$ this term should be finite. Choice of the small parameter ϵ is closely connected with that choice of parameter δ in the theorem below.

Theorem 2.8. *Assume that the class of probability distribution functions $\{B_1(x)\}$ is defined according to the above convention and satisfies (2.34), $\rho_1 = 1 - \delta$, $\delta > 0$, and $L\delta \rightarrow C > 0$, as $\delta \rightarrow 0$ and $L \rightarrow \infty$. Assume that $\rho_{1,3} = \rho_{1,3}(L)$ is a bounded sequence, and there exists $\widetilde{\rho}_{1,2} = \lim_{L \rightarrow \infty} \rho_{1,2}(L)$. Then,*

$$(2.35) \quad q_{L-j} = \frac{2\delta}{\widetilde{\rho}_{1,2}} \cdot \frac{1}{e^{2C/\widetilde{\rho}_{1,2}} - 1} \left(1 + \frac{2\delta}{\widetilde{\rho}_{1,2}}\right)^j [1 + o(1)]$$

for any $j \geq 0$.

Proof. We start the proof from the following auxiliary result, similar to that used to prove Theorem 2.6.

Lemma 2.9. *Under the assumptions of Theorem 2.8, the following asymptotic representation holds*

$$(2.36) \quad \tau = 1 + \frac{2\delta}{\widetilde{\rho}_{1,2}} + O(\delta^2).$$

Proof of Lemma 2.9. The proof of this lemma is completely similar to the proof of the result in [12], p.326 mentioned in the proof of Theorem 2.6 (see relation (2.23)). According to the above convention, the equation $z = \widehat{B}(\lambda - \lambda z)$ has a unique solution $\tau > 1$. Clearly, the root approaches 1 as δ vanishes, because otherwise we would have a contradiction with Corollary 2.2. Therefore, by the Taylor expansion of this equation around the point $z = 1$, we have

$$z = 1 - (1 - \delta)(1 - z) + \frac{\widetilde{\rho}_{1,2}}{2}(1 - z)^2 + O(1 - z)^3.$$

Ignoring the last term $O(1 - z)^3$ of this expansion, we have the quadratic equation, and the two solutions of the equation are $z = 1$ and $z = 1 + \frac{2\delta}{\widetilde{\rho}_{1,2}}$. Therefore,

$$\tau = 1 + \frac{2\delta}{\widetilde{\rho}_{1,2}} + O(\delta^2),$$

and the lemma is therefore proved. $\square$

Let us continue the proof of the theorem. From (2.33) and (2.36) of Lemma 2.9 we have:

$$(2.37) \quad q_{L-j} = q_L \left(1 + \frac{2\delta}{\widetilde{\rho}_{1,2}}\right)^j [1 + o(1)].$$

Taking into account that

$$\begin{aligned} \sum_{j=0}^{L-1} \left(1 + \frac{2\delta}{\widetilde{\rho}_{1,2}}\right)^j &= \frac{\widetilde{\rho}_{1,2}}{2\delta} \left[\left(1 + \frac{2\delta}{\widetilde{\rho}_{1,2}}\right)^L - 1 \right] \\ &= \frac{\widetilde{\rho}_{1,2}}{2\delta} \left(e^{2C/\widetilde{\rho}_{1,2}} - 1 \right), \end{aligned}$$

from the normalization condition $p_1 + p_2 + \sum_{j=1}^L q_j = 1$ and the fact that the both p_1 and p_2 have the order $O(\delta)$, we obtain:

$$(2.38) \quad q_L = \frac{2\delta}{\tilde{\rho}_{1,2}} \cdot \frac{1}{e^{2C/\tilde{\rho}_{1,2}} - 1} [1 + o(1)].$$

The desired statement of the theorem follows from (2.38). $\square$

Using the result of [5] we have the following.

Proposition 2.10. *Under the assumptions of Theorem 2.8 denote the objective function J by J^{lower} . We have the following representation*

$$(2.39) \quad J^{lower} = C \left[j_1 e^{\tilde{\rho}_{1,2}/2C} + j_2 \frac{\rho_2}{1 - \rho_2} \left(e^{\tilde{\rho}_{1,2}/2C} - 1 \right) \right] + c^{lower},$$

where

$$(2.40) \quad c^{lower} = \frac{2C}{\tilde{\rho}_{1,2}} \cdot \frac{1}{e^{2C/\tilde{\rho}_{1,2}} - 1} \lim_{L \rightarrow \infty} \frac{1}{L} \hat{C}_L \left(1 + \frac{2C}{\tilde{\rho}_{1,2}L} \right),$$

and $\hat{C}(z) = \sum_{j=0}^{L-1} c_{L-j} z^j$ is a backward generating cost function.

Proof. The representation for the term

$$C \left[j_1 e^{\tilde{\rho}_{1,2}/2C} + j_2 \frac{\rho_2}{1 - \rho_2} \left(e^{\tilde{\rho}_{1,2}/2C} - 1 \right) \right]$$

is known from [5]. The new term in (2.39), which takes into account the water costs, is c^{lower} . Keeping in mind representation (2.35), for this term we obtain:

$$c^{lower} = \lim_{L \rightarrow \infty} \sum_{j=0}^{L-1} q_{L-j} c_{L-j} = \lim_{L \rightarrow \infty} \sum_{j=0}^{L-1} c_{L-j} \cdot \frac{2\delta L}{\tilde{\rho}_{1,2}L} \cdot \frac{1}{e^{2C/\tilde{\rho}_{1,2}} - 1} \left(1 + \frac{2\delta L}{\tilde{\rho}_{1,2}L} \right)^j,$$

and representation (2.40) follows. $\square$

3. SOLUTION OF THE CONTROL PROBLEM AND ITS PROPERTIES

In this section we discuss the solution to the control problem and study its properties. The functionals J^{upper} and J^{lower} are given by (2.27) and (2.39). The last terms of these functionals are

$$(3.1) \quad c^{upper} = \frac{2C}{\tilde{\rho}_{1,2}} \cdot \frac{e^{2C/\tilde{\rho}_{1,2}}}{e^{2C/\tilde{\rho}_{1,2}} - 1} \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} c_{L-j} \left(1 - \frac{2\delta}{\tilde{\rho}_{1,2}} \right)^j,$$

and

$$(3.2) \quad c^{lower} = \frac{2C}{\tilde{\rho}_{1,2}} \cdot \frac{1}{e^{2C/\tilde{\rho}_{1,2}} - 1} \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} c_{L-j} \left(1 + \frac{2\delta}{\tilde{\rho}_{1,2}} \right)^j$$

correspondingly, where $C = \lim_{L \rightarrow \infty} L\delta$.

Now we give other representations for these terms. Denote

$$(3.3) \quad \psi(C) = \lim_{L \rightarrow \infty} \frac{\sum_{j=0}^{L-1} c_{L-j} \left(1 - \frac{2C}{\tilde{\rho}_{1,2}L} \right)^j}{\sum_{j=0}^{L-1} \left(1 - \frac{2C}{\tilde{\rho}_{1,2}L} \right)^j},$$

and

$$(3.4) \quad \eta(C) = \lim_{L \rightarrow \infty} \frac{\sum_{j=0}^{L-1} c_{L-j} \left(1 + \frac{2C}{\tilde{\rho}_{1,2}L}\right)^j}{\sum_{j=0}^{L-1} \left(1 + \frac{2C}{\tilde{\rho}_{1,2}L}\right)^j}.$$

Since $\{c_i\}$ is a bounded sequence, then the both limits of (3.3) and (3.4) do exist.

We have the following lemma.

Lemma 3.1. *We have*

$$(3.5) \quad c^{upper} = \psi(C),$$

and

$$(3.6) \quad c^{lower} = \eta(C).$$

Proof. From (3.3) and (3.4) we correspondingly have the representations

$$(3.7) \quad \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} c_{L-j} \left(1 - \frac{2C}{\tilde{\rho}_{1,2}L}\right)^j = \psi(C) \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} \left(1 - \frac{2C}{\tilde{\rho}_{1,2}L}\right)^j,$$

and

$$(3.8) \quad \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} c_{L-j} \left(1 + \frac{2C}{\tilde{\rho}_{1,2}L}\right)^j = \eta(C) \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} \left(1 + \frac{2C}{\tilde{\rho}_{1,2}L}\right)^j.$$

The desired results follow by direct transformations of the corresponding right-hand sides of (3.7) and (3.8).

Prove (3.5). For the right-hand side of (3.7) we have

$$(3.9) \quad \psi(C) \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} \left(1 - \frac{2C}{\tilde{\rho}_{1,2}L}\right)^j = \psi(C) \left(1 - e^{-2C/\tilde{\rho}_{1,2}}\right) \frac{\tilde{\rho}_{1,2}}{2C}.$$

On the other hand, from (3.1) we have

$$(3.10) \quad c^{upper} \left(1 - e^{-2C/\tilde{\rho}_{1,2}}\right) \frac{\tilde{\rho}_{1,2}}{2C} = \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} c_{L-j} \left(1 - \frac{2C}{\tilde{\rho}_{1,2}L}\right)^j.$$

Hence, from (3.7), (3.9) and (3.10) we obtain (3.3). The proof of (3.6) is completely analogous and uses representations (3.2) and (3.8). $\square$

The next lemma establishes the main properties of functions $\psi(C)$ and $\eta(C)$.

Lemma 3.2. *The function $\psi(C)$ is a non-increasing function, and its maximum is $\psi(0) = c^*$. The function $\eta(C)$ is an increasing function, and its minimum is $\eta(0) = c^*$. (Recall that $c^* = \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{i=1}^L c_i$ is defined in Proposition 2.4.)*

Proof. Let us first prove that $\psi(0) = c^*$ is a maximum of $\psi(C)$. For this purpose we use the following well-known inequality (e.g. Hardy, Littlewood and Polya [7] or Marschall and Olkin [8]). Let $\{a_n\}$ and $\{b_n\}$ be arbitrary sequences, one of them is increasing and another decreasing. Then for any finite sum we have

$$(3.11) \quad \sum_{n=1}^l a_n b_n \leq \frac{1}{l} \sum_{n=1}^l a_n \sum_{n=1}^l b_n.$$

Applying inequality (3.11) to finite sums of the left-hand side of (3.7) and passing to limit as $L \rightarrow \infty$, we have

$$\begin{aligned}
 (3.12) \quad & \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} c_{L-j} \left(1 - \frac{2C}{\tilde{\rho}_{1,2}L}\right)^j \\
 & \leq \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} c_{L-j} \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} \left(1 - \frac{2C}{\tilde{\rho}_{1,2}L}\right)^j \\
 & = \psi(0) \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} \left(1 - \frac{2C}{\tilde{\rho}_{1,2}L}\right)^j.
 \end{aligned}$$

Then, comparing (3.7) with (3.12) enables us to conclude,

$$\psi(0) = c^* \geq \psi(C),$$

i.e. $\psi(0) = c^*$ is the maximum value of $\psi(C)$.

Prove now, that $\psi(C)$ is a non-increasing function, i.e. for any nonnegative $C_1 \leq C$ we have $\psi(C) \leq \psi(C_1)$.

To prove this note, that for small positive δ_1 and δ_2 we have $(1-\delta_1-\delta_2) = (1-\delta_1)(1-\delta_2) + O(\delta_1\delta_2)$. Using this idea, one can prove the monotonicity of $\psi(C)$ by replacing

$$\left(1 - \frac{2C}{\tilde{\rho}_{1,2}L}\right) = \left(1 - \frac{2C_1}{\tilde{\rho}_{1,2}L}\right) \left(1 - \frac{2C - 2C_1}{\tilde{\rho}_{1,2}L}\right) + O\left(\frac{1}{L^2}\right), \quad C > C_1$$

in the above asymptotic relations for large L . Indeed, notice that

$$\begin{aligned}
 (3.13) \quad & \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} \left(1 - \frac{2C_1}{\tilde{\rho}_{1,2}L}\right)^j \left(1 - \frac{2C - 2C_1}{\tilde{\rho}_{1,2}L}\right)^j \\
 & = \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} \left(1 - \frac{2C_1}{\tilde{\rho}_{1,2}L}\right)^j \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} \left(1 - \frac{2C - 2C_1}{\tilde{\rho}_{1,2}L}\right)^j
 \end{aligned}$$

Therefore, for any non-decreasing sequence a_j

$$\begin{aligned}
 (3.14) \quad & \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} a_j \left(1 - \frac{2C_1}{\tilde{\rho}_{1,2}L}\right)^j \left(1 - \frac{2C - 2C_1}{\tilde{\rho}_{1,2}L}\right)^j \\
 & \leq \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} a_j \left(1 - \frac{2C_1}{\tilde{\rho}_{1,2}L}\right)^j \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} \left(1 - \frac{2C - 2C_1}{\tilde{\rho}_{1,2}L}\right)^j
 \end{aligned}$$

Indeed, assume for contrary that

$$\begin{aligned}
 (3.15) \quad & \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} a_j \left(1 - \frac{2C_1}{\tilde{\rho}_{1,2}L}\right)^j \left(1 - \frac{2C - 2C_1}{\tilde{\rho}_{1,2}L}\right)^j \\
 & > \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} a_j \left(1 - \frac{2C_1}{\tilde{\rho}_{1,2}L}\right)^j \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} \left(1 - \frac{2C - 2C_1}{\tilde{\rho}_{1,2}L}\right)^j
 \end{aligned}$$

Then, applying the inequality (3.11) to the right-hand side of (3.15), we obtain:

$$\begin{aligned}
 & \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} a_j \left(1 - \frac{2C_1}{\tilde{\rho}_{1,2}L}\right)^j \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} \left(1 - \frac{2C - 2C_1}{\tilde{\rho}_{1,2}L}\right)^j \\
 (3.16) \quad & \leq \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} a_j \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} \left(1 - \frac{2C_1}{\tilde{\rho}_{1,2}L}\right)^j \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} \left(1 - \frac{2C - 2C_1}{\tilde{\rho}_{1,2}L}\right)^j \\
 & = \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} a_j \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} \left(1 - \frac{2C}{\tilde{\rho}_{1,2}L}\right)^j.
 \end{aligned}$$

Comparison of the last obtained term with the left-hand side of (3.15) enables us to write:

$$\lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} a_j \left(1 - \frac{2C}{\tilde{\rho}_{1,2}L}\right)^j > \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} a_j \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} \left(1 - \frac{2C}{\tilde{\rho}_{1,2}L}\right)^j.$$

The contradiction with the basic inequality (3.11) proves (3.14).

Taking into account (3.13) and (3.14) the extended version of (3.12) after application (3.11) now looks

$$\begin{aligned}
 & \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} c_{L-j} \left(1 - \frac{2C}{\tilde{\rho}_{1,2}L}\right)^j \\
 & = \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} c_{L-j} \left(1 - \frac{2C_1}{\tilde{\rho}_{1,2}L}\right)^j \left(1 - \frac{2C - 2C_1}{\tilde{\rho}_{1,2}L}\right)^j \\
 (3.17) \quad & \leq \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} c_{L-j} \left(1 - \frac{2C_1}{\tilde{\rho}_{1,2}L}\right)^j \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} \left(1 - \frac{2C - 2C_1}{\tilde{\rho}_{1,2}L}\right)^j \\
 & = \psi(C_1) \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} \left(1 - \frac{2C_1}{\tilde{\rho}_{1,2}L}\right)^j \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} \left(1 - \frac{2C - 2C_1}{\tilde{\rho}_{1,2}L}\right)^j
 \end{aligned}$$

On the other hand, the right-hand side of (3.7) can be rewritten

$$\begin{aligned}
 & \psi(C) \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} \left(1 - \frac{2C}{\tilde{\rho}_{1,2}L}\right)^j \\
 (3.18) \quad & = \psi(C) \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} \left(1 - \frac{2C_1}{\tilde{\rho}_{1,2}L}\right)^j \left(1 - \frac{2C - 2C_1}{\tilde{\rho}_{1,2}L}\right)^j \\
 & = \psi(C) \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} \left(1 - \frac{2C_1}{\tilde{\rho}_{1,2}L}\right)^j \lim_{L \rightarrow \infty} \frac{1}{L} \sum_{j=0}^{L-1} \left(1 - \frac{2C - 2C_1}{\tilde{\rho}_{1,2}L}\right)^j.
 \end{aligned}$$

The last equality in (3.18) is an application of (3.13). From (3.17) and (3.18) we finally obtain $\psi(C_1) \geq \psi(C)$ for any positive $C_1 \leq C$. The first statement of Lemma 3.2 is proved. The proof of the second statement of this lemma is similar. $\square$

We are ready now to formulate and prove a main theorem on optimal control of the dam model considered in the present paper.

Theorem 3.3. *The solution to the control problem is $\rho_1 = 1$ if and only if the minimum of the both functionals J^{upper} and J^{lower} is achieved for $C = 0$. In this case the minimum of the functional J is given by (2.19). Otherwise, there can be one of the following two cases for the solution to the control problem.*

(1) *If the minimum of J^{upper} is achieved for $C = 0$, then the minimum of J^{lower} must be achieved for some positive value $C = \underline{C}$. Then the solution to the control problem is achieved for $\rho_1 = 1 - \delta$, where $\delta > 0$ vanishes such that $\delta L \rightarrow \underline{C}$ as $L \rightarrow \infty$.*

(2) *If the minimum of J^{lower} is achieved for $C = 0$, then the minimum of J^{upper} must be achieved for some positive value $C = \overline{C}$. Then the solution to the control problem is achieved for $\rho_1 = 1 + \delta$, where $\delta > 0$ vanishes such that $\delta L \rightarrow \overline{C}$ as $L \rightarrow \infty$.*

The minimum of the functionals J^{upper} or J^{lower} is defined by their differentiating and then equating these derivatives to zero.

Proof. Note first, that there is a unique solution of the control problem considered in this paper. Indeed, in the case where the water costs are not taken into account, the existence of a unique solution of the control problem follows from the main result of [5]. In the case of the dam model of this paper the only difference is in presence of the functions c^{upper} and c^{lower} in the functionals J^{upper} and J^{lower} correspondingly. According to Lemma 3.1 $c^{upper} = \psi(C)$ and $c^{lower} = \eta(C)$, and according to Lemma 3.2 the function $\psi(C)$ is non-increasing in C , while the function $\eta(C)$ is a non-decreasing in C , and $\psi(0) = \eta(0) = c^*$. Therefore, there is a unique solution of the control problem considered in the present paper as well.

In the case where the both minima of J^{upper} and J^{lower} are achieved in $C = 0$, then each of these minima is equal to the right-hand side of (2.19), and the minimum of the objective function J is achieved for $\rho_1 = 1$.

If the minimum of J^{lower} is achieved for $C = \underline{C}$, then J^{lower} is less than the right-hand side of (2.19), and, because of uniqueness of the solution, the minimum of J^{upper} must be achieved for $C = 0$. In this case the minimum of the objective function J is achieved for $\rho_1 = 1 - \delta$, $\delta > 0$ vanishes as $L \rightarrow \infty$, and $L\delta \rightarrow \underline{C}$.

In the opposite case, if the minimum of J^{upper} is achieved for $C = \overline{C}$, then J^{upper} is less than the right-hand side of (2.19), and the minimum of J^{lower} must be achieved for $C = 0$. In this case the minimum of objective function J is achieved for $\rho_1 = 1 + \delta$, $\delta > 0$ vanishes as $L \rightarrow \infty$, and $L\delta \rightarrow \overline{C}$. $\square$

From Theorem 3.3 we have the following evident property of the optimal control.

Corollary 3.4. *The solution of the control problem can be $\rho_1 = 1$ only in the case $j_1 \leq j_2 \frac{\rho_2}{1-\rho_2}$. Specifically, the equality is achieved only for $c_i \equiv c$, $i = 1, 2, \dots, L$, where c is an any positive constant.*

Although Corollary 3.4 provides a partial answer to the question 2 posed in the introduction, the answer is not satisfactory, because it is an evident extension of the result of [5]. A more constructive answer the question 2 of the introduction is obtained for the special case considered in the next section.

4. EXAMPLE OF LINEAR COSTS

In this section we study an example related to the case of linear costs.

Assume that c_1 and $c_L < c_1$ are given. Then the assumption that the costs are linear means, that

$$(4.1) \quad c_i = c_1 - \frac{i-1}{L-1}(c_1 - c_L), \quad i = 1, 2, \dots, L.$$

It is assumed that as L is changed, the costs are recalculated as follows. The first and last values of the cost c_1 and c_L remains the same. Other costs in the intermediate points are recalculated according to (4.1).

Therefore, to avoid confusing with the appearance of the index L for the fixed (unchangeable) values of cost c_1 and c_L , we use the other notation: $c_1 = \bar{c}$ and $c_L = \underline{c}$. Then (4.1) has the form

$$(4.2) \quad c_i = \bar{c} - \frac{i-1}{L-1}(\bar{c} - \underline{c}), \quad i = 1, 2, \dots, L.$$

In the following we shall also use the inverse form of (4.2). Namely,

$$(4.3) \quad c_{L-i} = \underline{c} + \frac{i}{L-1}(\bar{c} - \underline{c}), \quad i = 0, 1, \dots, L-1.$$

Apparently,

$$(4.4) \quad c^* = \frac{\bar{c} + \underline{c}}{2}.$$

For c^{upper} we have

$$(4.5) \quad \begin{aligned} c^{upper} = \psi(C) &= \lim_{L \rightarrow \infty} \frac{\sum_{j=0}^{L-1} \left(\underline{c} + \frac{j}{L-1}(\bar{c} - \underline{c}) \right) \left(1 - \frac{2C}{\tilde{\rho}_{1,2}L} \right)^j}{\sum_{j=0}^{L-1} \left(1 - \frac{2C}{\tilde{\rho}_{1,2}L} \right)^j} \\ &= \underline{c} + (\bar{c} - \underline{c}) \lim_{L \rightarrow \infty} \frac{1}{L-1} \cdot \frac{\sum_{j=0}^{L-1} j \left(1 - \frac{2C}{\tilde{\rho}_{1,2}L} \right)^j}{\sum_{j=0}^{L-1} \left(1 - \frac{2C}{\tilde{\rho}_{1,2}L} \right)^j} \\ &= \underline{c} + (\bar{c} - \underline{c}) \frac{\tilde{\rho}_{1,2}}{2C} \cdot \frac{-\frac{2C}{\tilde{\rho}_{1,2}} + e^{2C/\tilde{\rho}_{1,2}} - 1}{e^{2C/\tilde{\rho}_{1,2}} - 1}. \end{aligned}$$

For example, as C converges to zero in (4.5), then c^{upper} converges to $\underline{c} + \frac{1}{2}(\bar{c} - \underline{c}) = c^*$. This is in agreement with the statement of Proposition 2.4.

In turn, for c^{lower} we have

$$\begin{aligned}
 c^{lower} = \eta(C) &= \lim_{L \rightarrow \infty} \frac{\sum_{j=0}^{L-1} \left(\underline{c} + \frac{j}{L-1} (\bar{c} - \underline{c}) \right) \left(1 + \frac{2C}{\tilde{\rho}_{1,2} L} \right)^j}{\sum_{j=0}^{L-1} \left(1 + \frac{2C}{\tilde{\rho}_{1,2} L} \right)^j} \\
 (4.6) \quad &= \underline{c} + (\bar{c} - \underline{c}) \lim_{L \rightarrow \infty} \frac{1}{L-1} \cdot \frac{\sum_{j=0}^{L-1} j \left(1 + \frac{2C}{\tilde{\rho}_{1,2} L} \right)^j}{\sum_{j=0}^{L-1} \left(1 + \frac{2C}{\tilde{\rho}_{1,2} L} \right)^j} \\
 &= \underline{c} + (\bar{c} - \underline{c}) \frac{\tilde{\rho}_{1,2}}{2C} \cdot \frac{\frac{2C}{\tilde{\rho}_{1,2}} - 1 + e^{-2C/\tilde{\rho}_{1,2}}}{1 - e^{-2C/\tilde{\rho}_{1,2}}}.
 \end{aligned}$$

Again, as C converges to zero in (4.6), then c^{lower} converges to $\underline{c} + \frac{1}{2}(\bar{c} - \underline{c}) = c^*$. Again, we arrive to an agreement with the statement of Proposition 2.4.

Let us now discuss the question 2 posed in the introduction. We cannot give the explicit solution because the calculations are very routine and cumbersome. However, we explain the way of the solution of this problem and find a numerical result.

Following Corollary 3.4, take first $j_1 = j_2 \frac{\rho_2}{1-\rho_2}$. Clearly, that for these relation between parameters j_1 and j_2 the minimum of J^{lower} must be achieved for $C = 0$, while the minimum of J^{upper} must be achieved for a positive C . Now, keeping j_1 fixed assume that j_2 increases. Then, the problem is to find the value for parameter j_2 such that the value C corresponding to the minimization problem of J^{upper} reaches the point 0.

In our example we take $j_1 = 1$, $\rho_2 = \frac{1}{2}$, $\underline{c} = 1$, $\bar{c} = 2$, $\tilde{\rho}_{1,2} = 1$. In the table below we outline some values j_2 and the corresponding value C for optimal solution of functional J^{upper} . It is seen from the table that the optimal value is achieved in the case $j_2 \approx 1.34$. Therefore, in the present example $j_1 = 1$ and $j_2 \approx 1.34$ lead to the optimal solution $\rho_1 = 1$.

ACKNOWLEDGEMENT

The author acknowledges with thanks the support of the Australian Research Council, grant # DP0771338.

REFERENCES

- [1] ABRAMOV, V.M. (1997). On a property of a refusals stream. *Journal of Applied Probability*, **34**, 800-805.
- [2] ABRAMOV, V.M. (2002). Asymptotic analysis of the GI/M/1/n loss system as n increases to infinity. *Annals of Operations Research*, **112**, 35-41.
- [3] ABRAMOV, V.M. (2004). Asymptotic behavior of the number of lost messages. *SIAM Journal on Applied Mathematics*, **64**, 746-761.
- [4] ABRAMOV, V.M. (2006). The effective bandwidth problem revisited. arXiv: math. 0604182.
- [5] ABRAMOV, V.M. (2007). Optimal control of a large dam. *Journal of Applied Probability*, **44**, 249-258.
- [6] FELLER, W. (1966). *An Introduction to Probability Theory and Its Applications*, vol. 2. John Wiley, New York.

Parameter	Argument of optimal value
j_2	C
1.06	0.200
1.07	0.190
1.08	0.182
1.09	0.174
1.10	0.165
1.11	0.156
1.12	0.149
1.13	0.140
1.14	0.134
1.15	0.126
1.16	0.120
1.17	0.112
1.18	0.104
1.19	0.096
1.20	0.090
1.25	0.055
1.30	0.022
1.33	0.010
1.34	0

TABLE 1. The values of parameter j_2 and corresponding arguments of optimal value C

- [7] HARDY, G.H., LITTLEWOOD, J.E. AND POLYA, G. (1952). *Inequalities*. Second edn. Cambridge Univ. Press, London.
- [8] MARSCHALL, A.W. AND OLKIN, I. (1979). *Inequalities: Theory of Majorization and Its Applications*. Academic Press, London.
- [9] POSTNIKOV, A.G. (1980). *Tauberian Theory and Its Application*. Proc. Steklov Math. Inst. (2) **144** (AMS Transl. from Russian.)
- [10] RIGHTER, R. (1999). A note on losses in M/GI/1/n queues. *Journal of Applied Probability*, **36**, 1241-1244.
- [11] ROSS, S.M. (2000). *Introduction to Probability Models*, Seventh edn, Academic Press, Burlington.
- [12] SUBHANKULOV, M.A. (1976). *Tauberian Theorems with Remainder*, Nauka, Moscow. (Russian.)
- [13] TAKÁCS, L. (1962). *Introduction to the Theory of Queues*. Oxford Univ. Press, New York/London.
- [14] TAKÁCS, L. (1967). *Combinatorial Methods in the Theory of Stochastic Processes*. John Wiley, New York.
- [15] WILLMOT, G.E. (1988). A note on the equilibrium M/G/1 queue length. *Journal of Applied Probability*, **25**, 228-231.
- [16] WOLFF, R.W. (2002). Losses per cycle in a single-server queue. *Journal of Applied Probability*, **39**, 905-909.

SCHOOL OF MATHEMATICAL SCIENCES, MONASH UNIVERSITY, BUILDING 28M, WELLINGTON ROAD, CLAYTON, VIC 3800, AUSTRALIA
E-mail address: vyacheslav.abramov@sci.monash.edu.au